

Poštovani,

Učesnici XXXIV konferencije *PROCESNA TEHNIKA I ENERGETIKA U POLJOPRIVREDI PTEP 2022*, mogu odabrati i mogućnost da u Zborniku sa konferencije štampaju rad u celini (M33 kategorija). Zbornik (Proceedings) će biti publikovan u elektronskoj verziji. Da bi to i ostvarili treba da pripreme svoj rukopis za publikovanje. Rad treba bude kratak, 3-5 strana. Radovi za zbornik biće recenzirani.

Radove za objavljivanje u Zborniku radova sa konferencije pisati pre ma s ledeće m up utstv:

Format: A4

Margine: top 20 mm, bottom 20 mm, left 20 mm, right 20 mm

Naslov rada: Times New Roman, font 14, bold, centriran

Autori i pripadnost instituciji, e-mail: Times New Roman, font 12, Italic, centrirano

Rad na **engleskom jeziku** treba da sadrži sledeće poglavlja: uvod, materijal i metod (materijal-stručni rad), rezultati i diskusija (diskusija-stručni rad), zaključak, zahvalnica i literatura. Rad treba da bude napisan na **3-5 strana**. Radovi obima preko 5 strana i koji se nisu pridržavali ovog neće biti publikovani u ovoj formi (u celini).

Naslovi slika i tabela treba da budu *Italic*.

Primer pisanja rada

EFFECT OF THE PARAMETERS OF OSMOTIC DRYING ON SOME PHYSICAL PROPERTIES OF QUINCE

Petar PETROVIĆ¹, Milica MILIĆ¹, Marko MARKOVIĆ²

¹*University of Novi Sad, Faculty of Agriculture, Novi Sad, Trg Dositeja Obradovića 8, Serbia*

²*University of Belgrade, Faculty of Agriculture, Nemanjina 6, Serbia*

Contact: mradojcin@polj.uns.ac.rs

ABSTRACT (100 do 150 reči)

Osmotic drying of quince is used as the first step in combined drying. The removal of moisture during osmotic drying ensures less time for convective drying. Osmotic treatment reduces water activity, provides microbiological stability....

INTRODUCTION

Quince is a fruit species that is highly appreciated for its aroma. It is typically used in the production of brandy and compotes, while it is rarely used in the fresh state. This is because of the hardness and firmness of the fruit (Babić et al, 2008). Dried quince is used.....

MATERIAL AND METHOD

A variety of quince Leskovačka was used in the experiment. On the basis of previous research results this variety is very suitable from the aspect of size and shape.....

RESULTS AND DISCUSSION

During osmotic drying the color parameter b^* , which means yellow color, changed during all 180 minutes. There is a constant increase in the value of this parameter. Somewhat larger....

CONCLUSION

The physical properties of food products are the first indicator of quality. Preserving these properties is one of the goals to be achieved during the...

ACKNOWLEDGEMENTS (navesti samo broj projekta, bez naziva projekta)

This paper is a result of the research within the project TR31058, 2011-2014, supported by the Ministry of Education, Science and Technology, Republic of Serbia.

REFERENCES (max 10 referenci)

Alvarez-Fernandez, Ana, Pilar, P., Abadiäa, J., Anunciacioän, A. (2003). Effects of Fe deficiency chlorosis on yield and fruit quality in peach (*Prunus persica* L. Batsch). *Journal of Agricultural and Food Chiemistry*, 51, 5738-5744.